

Stroke in Minnesota At-A-Glance - 2007

Mortality

| | | Number of Deaths (2007) | Crude Death Rate per 100,000 | Age-Adjusted Death Rate per 100,000 |
|----------------|---|----------------------------|---------------------------------|--|
| Total | | 2081 | 40.0 | 36.2 |
| Race/Ethnicity | White | 1980 | 42.3 | 35.4 |
| | African-American/African American Indian | 47 | 18.3 | 52.3 |
| | Asian | 16 | - | - |
| | Hispanic | 28 | 14.3 | 38.7 |
| | | 23 | 11.2 | 37.5 |
| Male | Total | 823 | 31.8 | 37.7 |
| | White | 769 | 33.1 | 36.6 |
| | African-American/African American Indian | 23 | 17.1 | 56.9 |
| | Asian | 9 | - | - |
| | Hispanic | 14 | - | - |
| | | 14 | - | - |
| Female | Total | 1258 | 48.2 | 34.5 |
| | White | 1211 | 51.4 | 33.9 |
| | African-American/African American Indian | 24 | 19.5 | 49.1 |
| | Asian | 7 | - | - |
| | Hispanic | 14 | - | - |
| | | 9 | - | - |
| Age | 0-34 | 14 | - | |
| | 35-44 | 22 | 3.0 | |
| | 45-54 | 82 | 10.2 | |
| | 55-64 | 113 | 20.2 | |
| | 65-74 | 227 | 71.7 | |
| | 75-84 | 586 | 272.9 | |
| | 85+ | 1037 | 988.9 | |

Note: Rates for cells with fewer than 20 respondents are not reported, as estimates are unstable.

Source: Minnesota Department of Health

Prevalence of Stroke

| | | Percent |
|----------------|--------------------------|---------|
| Overall | | 2.2 |
| Sex | Male | 2.1 |
| | Female | 2.3 |
| Race/Ethnicity | White | 2.3 |
| | African-American/African | 1.3 |
| | Hispanic | 0.9 |
| | Other | 0.9 |
| | Multiracial | - |
| Age | 18-24 | - |
| | 25-34 | - |
| | 35-44 | - |
| | 45-54 | - |
| | 55-64 | 2.5 |
| | 65+ | 8.6 |
| Income | <\$15,000 | 9.2 |
| | \$15,000-24,999 | 4.4 |
| | \$25,000-34,999 | 3.4 |
| | \$35,000-49,999 | 2.4 |
| | \$50,000 + | 0.9 |
| Education | Less than HS | - |
| | HS or GED | 3.3 |
| | Some post HS | 1.9 |
| | College graduate | 1.2 |

Source: Minnesota Behavioral Risk Factor Surveillance System Survey 2008

Note: Percents for cells with fewer than 20 respondents are not reported, as estimates are unstable.